

(19)



(11)

EP 2 592 085 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

15.05.2013 Bulletin 2013/20

(51) Int Cl.:

C07D 513/04 (2006.01)

(21) Application number: **11188961.4**

(22) Date of filing: **14.11.2011**

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

Designated Extension States:

BA ME

(71) Applicant: **King Saud University**

11421 Riyadh (SA)

(72) Inventors:

- **Al-Rashood, Sara T. A.**
Riyadh 11333 (SA)

• **EI-Subbagh, Hussein I.**

11421 Riyadh (SA)

• **Hassan, Ghada S.**

Riyadh 11333 (SA)

• **EI-Taheer, Kamal E. H.**

Riyadh 11333 (SA)

• **Al-Omar, Mohamed A.**

11421 Riyadh (SA)

(74) Representative: **Goddar, Heinz J.**

Boehmert & Boehmert

Pettenkofenstrasse 20-22

80336 München (DE)

(54) **Thiazolo[3,2-a][1,3]diazepine derivatives and pharmaceutical compositions containing the same as anticonvulsant agents and method for their preparation**

(57) The present invention relates to compounds of the formula 1, a pharmaceutical composition containing the compounds, a method for their preparation as well as the use of the compounds as anticonvulsant agents.

EP 2 592 085 A1

Description

[0001] The present invention relates to compounds useful as anticonvulsant agents, pharmaceutical composition containing the compounds as well as a method for their preparation.

[0002] Epilepsy is one of the most common neurological disorders, which is characterized by seizures that take various forms and result from episodic neural discharges. The forms of seizures depend on the part of the brain affected. Epilepsy affects 0.5-1% of the population. Although there is no recognizable cause which can be identified, patient may develop seizure after brain trauma, infection or tumor growth, or other kinds of neurological diseases, including various inherited neurological syndromes. Epilepsy is treated mainly with drugs, though brain surgery may be used for severe cases. Current antiepileptic drugs are effective in controlling seizures in about 70% of patients, but their use is often limited by side-effects (Bordie M.J., Lancet. 1990, 336, 350-354; Shoravon S.D., Lancet. 1990, 336, 93-96; Meldrum B.S. et al. In New Anticonvulsant Drugs. Current Problems in Epilepsy 4; John Libby: London, 1986. Flaherty P.T. et al., J. Med. Chem. 1996, 39, 1509-1513).


[0003] GABA (γ -aminobutyric acid) is the principal inhibitory transmitter of many CNS pathways regulating numerous neurological functions including convulsions, anxiety and sleep activity. GABA acts on the GABA_A chloride ion channel. Molecular biology studies have demonstrated that several different receptor subunits combine to form the GABA_A receptor complex with a number of receptor subtypes such as benzodiazepine receptor (BzR). Ligands acting at the BzR allosteric binding site of the GABA_A receptor modulate the action of GABA on chloride ion flux. They show a wide variety of pharmacological actions ranging from full agonistic effects such as sedative/hypnotic, anxiolytic and anticonvulsant activities; to inverse agonistic effects such as anxiogenic, and proconvulsant activities. Several classes of chemical compounds such as benzodiazepines, steroids and barbiturates bind to the GABA_A chloride ion channel complex. The azepine derivatives such as imidazo[2,1-b]thiazepine exhibited affinity for BzR. Their profile of action was characterized as partially agonistic at the BzRs on the basis of the GABA shift.

[0004] The α -Amino-3-hydroxy-5-methyl-4-isoxazole propionate receptors (AMPA_Rs) are glutamate-gated ion channels which mediate the majority of fast excitatory synaptic transmissions in the mammalian brain. Hyperactivation of AMPARs is implicated in a broad range of acute and chronic neurodegenerative and neuropsychiatric conditions such as epilepsy. Thus, the modulation of AMPARs as potential targets for therapeutic intervention in many neurological disorders is one of the goals in neuropharmacology (Dingledine R. et al. Pharmacol. Rev. 1999, 51, 7-61; Lees G. J., Drugs 2000, 59, 33-78). Consequently, several antagonists of AMPAR such as talampanel (Rogawski M. A. Trends Pharmacol. Sci. 1993, 14, 325-331) and CFM-2 (Quartarone S., et al. IL Farmaco 2004, 59, 353-358) have been reported and showed promising therapeutic potential for the prevention and treatment of epilepsy.

[0005] DE 103 20 732 relates to 3-carboxylic acid diazepine derivatives and its use as rapid-acting anesthetics. The publication of Kadi et al. analyst, 2011, 136, 591-597 relates to liquid chromatographic high-throughput analysis of a new ultra short acting hypnotic 3-carboxylic acid diazepine derivative and its metabolic in mice serum using a monolithic silica column.

[0006] It is an object of the present invention to provide an anticonvulsant agent which overcomes the drawbacks of the prior art. Especially a compound shall be provided exhibiting potent in vivo anticonvulsant activity. Additionally, a pharmaceutical composition containing such anticonvulsant agent shall be provided, as well as a method for its preparation

[0007] This object is achieved by a compound according to formula 1:



1

wherein R_1 , R_2 , R_3 , R_4 and R_5 are each independently selected from the group consisting of hydrogen, halogen, C_1 - C_{20} -alkyl, C_1 - C_{20} -haloalkyl, C_1 - C_{20} -alkoxy, C_1 - C_{20} -haloalkoxy, aryl, heteroaryl, mercapto, alkylthio, amino, alkylamino, or wherein R_2 , R_3 and R_4 are a member of a ring system.

[0008] In an preferred embodiment, R_2 is selected from the group consisting of hydrogen, mercapto and C_1 - C_{20} -alkyl, preferably methyl, ethyl, propyl, isopropyl or butyl, halogen or amino.

[0009] Even preferred, R_3 is hydrogen or is taken together with R_4 to form an, optionally substituted, alicyclic, aryl, or heteroaryl ring system.


[0010] Preferably, R_1 , R_2 , R_3 , R_4 and R_5 are independently selected from aryl, more preferably phenyl and naphthyl,

or heteroaryl, more preferably furyl, pyrrolyl, thienyl, imidazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, benzothiazolyl, and oxadiazolyl.

[0011] Preferred, the compound is present in form of its addition salt, preferably hydrochloride, hydrobromide, phosphate, nitrate, acetate, malate, succinate, fumarate, tartrate, salicylate, sorbate, lactate, p-toluene sulphate, or naphthalene-1,5-disulfonate salts.

[0012] The object is further achieved by a method for preparing an inventive compound, comprising: reacting a compound according to formula 2 with a compound according to formula 3 to prepare a compound according to formula 4, and reacting the compound of formula 4 to result in a compound of formula 1, as given in scheme 1:

Scheme 1



with R_1 , R_2 , R_3 , R_4 and R_5 are each independently selected from the group consisting of hydrogen, halogen, C_1 - C_{20} -alkyl, C_1 - C_{20} -haloalkyl, C_1 - C_{20} -alkoxyl, C_1 - C_{20} -haloalkoxy, aryl, heteroaryl, mercapto, alkylthio, amino, alkylamino, or wherein R_2 , R_3 and R_4 are a member of a ring system.

[0013] Preferably, the reaction of compound 2 and compound 3 is in the presence of potassium carbonate in a solvent, preferably toluene, ethyl benzene, o-xylene, m-xylene, p-xylene, octane, nonane and/or isopropyl benzene.

[0014] Even preferred, compound 4 is cyclized in the presence of at least one secondary amine, preferably diethyl amine, pyrrolidine, morpholine, piperidine and/or N-methylpiperazine, in a solvent, preferably toluene, ethylbenzene, o-xylene, m-xylene, p-xylene and/or isopropylbenzene.

[0015] The object is also achieved by a pharmaceutical composition comprising at least one of the inventive compounds and a pharmaceutically acceptable carrier or excipient.

[0016] Finally, the object is achieved by the use of the inventive compound or the inventive pharmaceutical composition as anticonvulsant agent for treatment of convulsion especially for treatment of epilepsy.

[0017] Surprisingly, it was found that the compounds according the present invention possess anticonvulsant activity, preferably superior over the activity known for the state of art. In particular, it was found that compounds according to the present invention exhibit an activity 2-4 fold higher than that of valproate sodium which is used as a standard anticonvulsant agent in treatment of epilepsy.


[0018] The term "alkyl" with regard to the definition of R_1 - R_5 in the compound according to formula 1 is to be understood to comprise linear and branched alkyls. The term "halo" shall comprise derivatives which are mono-, di-, tri- or poly-halosubstituted.

[0019] If possible, all substituents R_1 - R_5 may be optionally further substituted, for example by halogen, amino, substituted amino, C_1 - C_{20} -alkyl, C_1 - C_{20} -haloalkyl, C_1 - C_{20} -alkoxy or C_1 - C_{20} -haloalkoxy, alkylamino, arylthio, heteroarylthio, arylamino or heteroarylamino, carboxylate, ethyl carboxylate.

[0020] Thiazolo[3,2-a][1,3]diazepine analogues could be obtained adopting published methods (Molina, P. et al, J.

Org. Chem. 1993, 58, 5264-5270; Imming, P. et al, Arch. Pharm. (Weinheim) 1995, 238, 207-215; Lehmann, J. El-Subbagh, H. I. El-Kashef, H. A. 2004, DE 103 20 732 A1; El-Subbagh, H.I. et al. Bioorg. & Med. Chem. let. 2008,18, 72-77). The compounds of the invention and their analogues (**1**) are synthesized according to an inventive method, Scheme 1. The proper 2-amino-4, 5-substituted-thiazoles (**2**) were acylated with the suitable acid chloride derivatives (**3**), where Y is chlorine or bromine, preferably bromine, and anhydrous potassium carbonate in a suitable solvent, such as, for example, toluene, ethylbenzene, o-, m-, and p-xylene, octane, nonane and isopropylbenzene, preferably toluene and ethylbenzene, at temperature ranging from room temperature to 150°C, preferably 100-120°C. The products **4** can be purified by silica gel and neutral alumina chromatography. Compounds of the formula **4** were cyclized preferably using secondary amines, such as for example, diethylamine, pyrrolidine, morpholine, piperidine, N-methylpiperazine, preferably pyrrolidine and piperidine, in a suitable solvent, such as for example toluene, ethylbenzene, o-, m-, and p-xylene, isopro-pylbenzene, preferably toluene, o-xylene at temperature ranging 100-180°C, preferably 120-130°C. The products **1** can be purified by silica gel and neutral alumina chromatography. Representative examples of such synthesis are shown in Examples 1-4 below.

Scheme 1



[0021] Adult male Swiss albino mice (22-28 g), approximately 10-week old, were used to conduct the anticonvulsant evaluation. They were housed in cages and kept at a temperature of $20 \pm 2^\circ C$ and a relative humidity of $55 \pm 5\%$ with a light-dark cycle of 12 h and fed with standard diet and water *ad libitum*. Compounds of the formula **1** have been dissolved in dimethylsulfoxide and administered at dose levels of 25, 50, 100 and 200 mg/kg intraperitoneal (i.p.). Sodium valproate was freshly dissolved in 0.9 % NaCl and used at dose level of 200 mg/kg (1.38 mmol/kg) i.p., 30 min after the test compounds as a positive control. A control group of mice was included and received intraperitoneal (i.p.) injections of the test compounds' vehicle. Pentylenetetrazole (PTZ) was freshly dissolved in 0.9 % NaCl and used at dose level of 100 mg/kg, i.p. This dose was found to be the minimum dose that induced 100% clonic convulsion. Picrotoxin (pic) was freshly dissolved in 1 ml 0.1N warmed HCl and the final volume was made up with 0.9 % NaCl. It has been used at dose level of 10 mg/kg i.p. and was given 30 min after the test compounds.

[0022] The anticonvulsant activity of compounds of the formula **1** in mice was measured using Pentylenetetrazole (PTZ) seizure threshold test (White, H.S. et al. In Antiepileptic Drugs, 4th ed.; Levy, R.H. et al., Eds, Raven: New York, 1995, pp 99-121). The active compounds of the formula **1** were further tested against maximal electroshock seizure (MES), and picrotoxin induced clonic convulsions, to explore the involvement of GABA-ergic receptors in their anticonvulsant activity. The effect of compounds of the formula **1** on motor performance was studied by chimney test (Dauge, V. et al. Neuropsychopharmacology 2001, 25, 690-98). Median effective (ED_{50}), and median sedative (TD_{50}) were calculated. Representative examples are shown in Table 1, Examples 5-9.

[0023] The acute toxicity determination (LD_{50}) for compounds of formula **1** was performed. Compounds were given

intraperitoneally (i.p.) in doses ranging from 0.1-5 mmol/kg. The animals were observed for up to 6 hours continuously and were then kept under observation for 72 hours. All behavioral changes and death during the observation periods were recorded. The percentage of death at each dose level was then calculated, and the LD₅₀ values were obtained (Ghosh, M., Fundamental of Experimental Pharmacology, Scientific Book Agency, Calcutta. 1984, pp 153-158, 187-189).
5 The protective index (PI) and therapeutic index (TI) of each compound were calculated. Representative examples are shown in Table 1, Examples 10-11.

[0024] Biological evaluation of the new compounds of the formula 1 of the invention revealed that the compounds possess a remarkable anticonvulsant activity. The obtained results clearly point to the discovery of a new group of anticonvulsant agents that induce their actions via interaction with GABA_A receptors. The safety of the new compounds of the formula 1 of the invention is comparable to that of valproate sodium. Therefore, compounds of the formula I of the invention have the potential use as anticonvulsants, with potency proved to be 2-4 fold more active than valproate sodium.

[0025] Compounds of the formula 1 of the invention, and their acid addition salts display anticonvulsant activity. The present invention includes pharmaceutical formulations which, in addition to non-toxic, inert pharmaceutically suitable excipients, contain one or more active compounds according to the invention, or which consist of one or more active compounds according to the invention, as well as processes for the preparation of these formulations.

[0026] The present invention also includes pharmaceutical formulations in dosage units. This means that the formulations are in the form of individual parts, for example tablets, dragees, capsules, pills, and ampoules, of which the content of active compound corresponds to a fraction or a multiple of an individual dose. The dosage units can contain, for example, 1, 2, 3 or 4 individual doses or 1/2, 1/3 or 1/4 of an individual dose. An individual dose preferably contains the amount of active compound which is given in one administration and which usually corresponds to a whole, a half, a third or a quarter of a daily dose.

[0027] By non-toxic, inert pharmaceutically suitable excipients there are to be understood solid, semi-solid or liquid diluents, fillers and formulations auxiliaries of every kind.

[0028] Tablets, dragees, capsules, pills, granules, solutions and sprays may be mentioned as preferred pharmaceutical formulations

[0029] Tablets, dragees, capsules and pills can contain the active compound or compounds alongside the customary excipients, such as (a) fillers and extenders, for example starches, lactose, sucrose, glucose, mannitol and silica, (b) binders, for example carboxymethylcellulose, alginates, gelatin and polyvinylpyrrolidone, (C) humectants, for example agar-agar, calcium carbonate and sodium bicarbonate, (e) solution retarders, for example paraffin, and (f) resorption accelerators, for example quaternary ammonium compounds, (g) wetting agents, for example cetyl alcohol and glycerol monostearate, (h) adsorbents for example kaolin and bentonite, and (i) lubricants, for example talc, calcium stearate and magnesium stearate and solid polyethylene glycols, or mixtures of the compounds listed under (a) to (i).

[0030] The tablets, dragees, capsules and pills can be provided with the customary coatings and shells, optionally containing pacifying agents, and can also be of such composition that they release the active compound or compounds only, or preferably, in a certain part of the intestinal tract, optionally in a delayed manner, examples of embedding compositions which can be used being polymeric substances and waxes.

[0031] The active compound or compounds, optionally together with one or more of the above mentioned excipients could also be in a micro-encapsulated form.

[0032] Solutions and emulsions for parenteral administration can contain, in addition to the active compound or compounds, the customary excipients, such as solvents, solubilizing agents and emulsifiers, for example water, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils, especially cotton seed oil, groundnut oil, maize germ oil, olive oil, castor oil and sesame oil, glycerol, glycerol-formal, tetrahydrofurfuryl alcohol, polyethylene glycol and fatty acid esters of sorbitol, or mixtures of these substances, in a sterile form which is isotonic with blood.

[0033] The therapeutically active compounds should preferably be present in the above-mentioned pharmaceutical formulations in a concentration of about 0.1 to 99.5, preferably of about 0.5 to 95% by weight of the total mixture.

[0034] The above-mentioned pharmaceutical formulations can also contain other pharmaceutical formulations and/or other pharmaceutical active compounds in addition to the active compounds according to the invention.

[0035] The above-mentioned pharmaceutical formulations are prepared in the customary manner according to known methods, for example by mixing the active compound or compounds with the excipient or excipients.

[0036] The present invention also includes the use of the active compounds according to the invention, and of pharmaceutical formulations which contain one or more active compounds according to the invention in human and veterinary medicine.


[0037] The actual dosage unit will be determined by such generally recognized factors as body weight of the patient and/or severity and type of pathological condition the patient might be suffering. With these considerations in mind, the dosage unit for a particular patient can be readily determined by the medical practitioner in accordance with the techniques known in the medical arts.

[0038] The precise instructions for pharmaceutical administration of the compounds and agents according to the invention necessarily depend on the requirements of the individual case, the nature of treatment, and of course the opinion of the treating physician.

5 **Example 1**

Ethyl 2-(4-chlorobutanamido)-4-methylthiazole-5-carboxylate.

[0039]




[0040] A mixture of ethyl 2-amino-4-methylthiazole-5-carboxylate (7.44 g, 0.04 mol), 4-chlorobutyl chloride (11.3 g, 0.08 mol), and potassium carbonate (5.5 g, 0.04 mol) in toluene (100 ml) was stirred at room temperature for 24 h. The toluene was then evaporated under reduced pressure. The residue was then quenched with water, stirred, and filtered. The solid obtained was washed, dried and recrystallized from chloroform to give the required product (8.7 g, 75% yield), mp 140-2°C, m/e 290.6, 100% (consistent with molecular formula C₁₁H₁₅ClN₂O₃S, calcd. 290.77). ¹H NMR (CDCl₃): δ 1.29 (t, 3H, J = 7.0 Hz, CH₃-CH₂-), 2.14 (t, 2H, J = 6.0 Hz, -CH₂-CH₂-CH₂-), 2.59-2.61 (m, 5H, CH₃- & -CH₂-CH₂-CH₂-), 3.57 (t, 2H, J = 6.0 Hz, -CH₂-CH₂-CH₂-), 4.24-4.28 (q, 2H, J = 7.0 Hz, CH₃-CH₂-), 9.92 (brs, 1H, NH). ¹³C NMR: δ 14.3, 17.3, 27.2, 32.8, 43.6, 61.1, 116.1, 155.6, 160.1, 162.6, 170.2.

Example 2

Ethyl 3-methyl-8-oxo-5,6,7,8-tetrahydro-thiazolo[3,2-a][1,3]diazepin-2-carboxylate (SGH-35).

[0041]




[0042] A mixture of ethyl 2-(4-chlorobutanamido)-4-methylthiazole-5-carboxylate (1.16 g, 0.004 mol) and diethylamine (0.8 ml, 0.008 mol) in toluene (50 ml) was heated under reflux for 3 h. The reaction mixture was cooled, poured into water and stirred. Toluene was separated, dried and evaporated to give a crude product which was purified by repeated silica gel column chromatography eluting with CHCl₃/Hexane (90:10 v/v). The solid obtained was washed, dried and recrystallized from EtOH/H₂O to give the required product (0.41 g, 40% yield); mp 95-7°C; m/e 254.6, 100% (consistent with molecular formula C₁₁H₁₄N₂O₃S, calcd. 254.31). IR (KBr, cm⁻¹): 3026 (CH₃), 2980 (CH₂), 1707, 1695 (2 CO). ¹H NMR (CDCl₃): δ 1.26 (t, 3H, J = 7.0 Hz, CH₃-CH₂-), 2.16-2.22 (m, 2H, J = 7.5 Hz, -CH₂-CH₂-CH₂-), 2.58 (s, 3H, CH₃-), 2.63 (t, 2H, J = 8.0 Hz, -CH₂-CH₂-CH₂-), 4.08 (t, 2H, J = 7.5 Hz, -CH₂-CH₂-CH₂-), 4.21-4.26 (q, 2H, J = 7.0 Hz, CH₃-CH₂-). ¹³C NMR: δ 14.4, 17.3, 18.1, 31.6, 47.6, 60.8, 116.00, 156.5, 158.4, 162.9, 174.0.

Example 3

N-(5-Bromo-4-methylthiazol-2-yl)-4-chlorobutanamide

[0043]




[0044] A mixture of 5-bromo-4-methylthiazol-2-amine (7.72 g, 0.04 mol), 4-chlorobutyl chloride (11.3 g, 0.08 mol), and potassium carbonate (5.5 g, 0.04 mol) in toluene (100 ml) was stirred at room temperature for 24 h. The toluene was then evaporated under reduced pressure. The residue was then quenched with water, stirred, and filtered. The solid obtained was washed, dried and recrystallized from chloroform to give the required product (7.5 g, 63% yield), mp 112-4°C, m/e 297.6, 25% (consistent with molecular formula C₈H₁₀BrClN₂OS, calcd. 297.6). ¹H NMR (DMSO-d₆): δ 2.02-2.08 (m, 2H, J = 7.0 Hz, -CH₂-CH₂-CH₂-), 2.61-2.65 (m, 5H, CH₃- & -CH₂-CH₂-CH₂-), 3.69 (t, 2H, J = 7.0 Hz, -CH₂-CH₂-CH₂-), 12.42 (brs, 1H, NH). ¹³C NMR: δ 14.7, 27.3, 32.0, 44.7, 62.9, 158.3, 159.1, 170.4.

Example 4

2-Bromo-3-methyl-6,7-dihydro-thiazolo[3,2-a][1,3]diazepin-8(5H)-one (SGH-40)

[0045]



[0046] A mixture of N-(5-bromo-4-methylthiazol-2-yl)-4-chlorobutanamide (1.19 g, 0.004 mol) and diethylamine (0.8 ml, 0.008 mol) in toluene (50 ml) was heated under reflux for 3 h. The reaction mixture was cooled, poured into water and stirred. Toluene was separated, dried and evaporated to give a crude product which was purified by repeated silica gel column chromatography eluting with CHCl₃/Hexane (90:10 v/v). The solid obtained was washed, dried and recrystallized EtOH/H₂O to give the required product (0.6 g, 57% yield), mp 165-7°C, m/e 261.7, 20% (consistent with molecular formula C₈H₉BrN₂OS, calcd. 261.14). IR (KBr, cm⁻¹): 3021 (CH₃), 2992 (CH₂), 1707 (CO). ¹H NMR (DMSO-d₆): δ 2.15-2.21 (m, 2H, J = 7.0 Hz, -CH₂-CH₂-CH₂-), 2.60-2.65 (m, 5H, CH₃-CH₂-CH₂-CH₂-), 4.04 (t, 2H, J = 7.0 Hz, -CH₂-CH₂-CH₂-). ¹³C NMR: δ 14.9, 17.7, 30.7, 47.6, 157.3, 159.0, 159.8, 173.8.

[0047] The NMR spectral data assignments of compounds of Example 1-4 are based on analysis of the ¹H, Attached Proton Test (APT), the Distortionless Enhancement Polarization Transfer (DEPT), correlated spectroscopy (COSY), Heteronuclear Multiple Quantum Coherence Spectroscopy (HMQC), NMR spectra for each compound

Example 5

Effect of SGH-35 and SGH-40 on Pentylene-tetrazole (PTZ)-induced convulsion

[0048] Pentylene-tetrazole (PTZ) seizure threshold test is used to evaluate the potential anticonvulsant activity of compounds of the formula 1 (White, H.S. et al. In Antiepileptic Drugs, 4th ed.; Levy, R.H. et al., Eds, Raven: New York, 1995, pp 99-121). Each compound was tested in 4 mice groups (Swiss albino mice, 25 g body weight) for each dose level (25, 50, 100 and 200 mg/kg) and was given i.p. 30 min later, animals were received PTZ (100 mg./kg, i.p.) and observed for 30 min. A single 5 seconds episode of clonic spasms was taken as a threshold seizure. SGH-35 and SGH-40 produced 100% protection against PTZ-induced seizures, Table 1.

Table 1: ED₅₀, TD₅₀, PI, LD₅₀ and TI for SGH-35 and SGH-40.

Compd ^a	Dose		% Protec-tion	ED ₅₀ ^d mg/kg	TD ₅₀ ^d mg/kg	PI ^e	LD ₅₀ mg/kg	TI ^f	
	mmol/kg	mg/kg							
SGH-35	0.78	200	PTZ	100	127±9.3	200±12.3	1.57	670	5.28
			MES ^b	100	90±5.7	200±12.3	2.22		
			Pic ^c	100	148±11.1	200±12.3	1.35		
SGH-40	0.38	100	PTZ	100	112±3.9	170±6.9	1.51	640	5.71
			MES ^b	100	78±4.1	170±6.9	2.17		

(continued)

Compd ^a	Dose		% Protec- tion	ED ₅₀ ^d mg/kg	TD ₅₀ ^d mg/kg	PI ^e	LD ₅₀ mg/kg	TI ^f
	mmol/kg	mg/kg						
		Pic ^c	100	129±6.1	170±6.9	1.31		

^a Each dose of selected compounds was tested using 4 animals and the percentage of animals protected was recorded and the anticonvulsant activity was calculated.

^b MES: Maximal electroshock test, was applied 30 min after the administration of each compound.

^c Pic: Picrotoxin (8 mg/kg, IP) was given 30 min after the administration of each compound.

^d ED₅₀ and TD₅₀: Median effective and median sedative doses, respectively.

^e PI: Protective index = TD₅₀/ED₅₀

^f TI: Therapeutic index = LD₅₀/ED₅₀.

Example 6**Effect of SGH-35 and SCH-40 against the maximal electroshock (MES)-induced convulsion**

[0049] The maximal electroshock (MES) seizure is one of the electrical test used to evaluate anticonvulsant activity (White, H.S. et al. In Antiepileptic Drugs, 4th ed.; Levy, R.H. et al., Eds, Raven New York, 1995. pp 99-121). MES that induced 100% maximal seizure was found to be 50 mA alternating current of 100Hz frequency for 0.2 seconds, using ECT UNIT (model number 7801, UGO Basile, Varese, Italy). Swiss albino mice (25 g body weight) were injected with 25, 50, 100 and 200 mg/kg of SGH-35 and SGH-40 i.p. 30 min later, mice were restrained by hand and subjected to electric shock through their ears, and released immediately following electrical stimulation, to permit observation of the maximal seizure. The maximal seizure typically consists of a short period of initial tonic flexion and a prolonged period of tonic extension (especially the hind limb). Protection was defined as complete absence of hind limb tonic extension (Table 1).

Example 7**Effect of SGH-35 and SGH-40 against picrotoxin (pic)-induced convulsion**

[0050] Swiss albino mice (25 g body weight) were injected with 25, 50, 100 and 200 mg/kg of **SGH-35** and **SGH-40** (4 mice, i.p.). 30 min later, animals were injected sc with Pic (10 mg/kg) and observed for 30 min and the percentage protection against convulsive seizures for each compound was calculated (Table I).

Example 8**Minimal neurotoxicity (chimney test) for SGH-35 and SGH-40**

[0051] The effect of SGH-35 and SGH-40 on motor performance was studied by chimney test (Dauge, V. et al. Neuropsychopharmacology 2001, 25, 690-98). At 15 and 25 min after i.p. administration of compounds SGH-35 and SGH-40, the inability of mice to climb up backwards in a glass tube of 25 cm length and 3 cm inner diameter within 30 sec was recorded and taken as a measure of neurological deficits. Normal mice climb up in 5-10 sec. The dose that impaired the motor coordination in 50% of the animals tested was labeled as the median toxic dose 50 (TD₅₀).

Example 9**Calculation of the anticonvulsant ED₅₀ values for SGH-35 and SGH-40**

[0052] In each of the chemical and the electrical methods described above, Swiss albino mice were divided into 4 groups (N = 4 animals). Each group was injected (i.p.) with one of the four selected doses that have proven to obtain a variable percentage of protection 20 minutes before the convulsants. The effective dose 50 (ED₅₀) i.e. the dose that protected half of the animals was then calculated using the log-probit method (White, H.S. et al. In Antiepileptic Drugs, 4th ed.; Levy, R.H. et al., Eds, Raven: New York, 1995, pp 99-121).

Example 10**Determination of LD₅₀ and dose-response curve:**

5 [0053] Initially the approximate LD₁₀₀ of each test compound was determined. Albino Swiss mice (25 g body weight) were then divided into 4 groups (6 mice each). Each group was then injected with one of 4 selected doses the highest being the LD₁₀₀ dose. The animals were then observed closely for 4 hours and then occasionally for 72 hours. The percentage of death in each group was then calculated. The LD₅₀ doses were then calculated using the Probit method (Ghosh; M., Fundamentals of Experimental Pharmacology, Scientific Book Agency, Calcutta. 1984, pp 153-158, 10 187-189).


Example 11**Calculation of the protective Index (PI) and Therapeutic Index for SGH-35 and SGH-40**

15 [0054] The protective index (PI) - i.e. the value that indicates the margin of safety and tolerance of the test compound was calculated by the formula: Protective index (PI) = TD₅₀/ED₅₀. The Therapeutic Index was calculated by the formula: Therapeutic index (TI) = LD₅₀/ED₅₀ (Löscher, W. et al. Epilepsy Res. 1991, 9, 1-10).

20 [0055] The features disclosed in the foregoing description and in the claims may, both separately and in any combination thereof, be material for realizing the invention in diverse forms thereof.

Claims

25 1. Compound according to formula 1:



wherein R₁, R₂, R₃, R₄ and R₅ are each independently selected from the group consisting of hydrogen, halogen, C₁-C₂₀-alkyl, C₁-C₂₀-haloalkyl, C₁-C₂₀-alkoxy, C₁-C₂₀-haloalkoxy, aryl, heteroaryl, mercapto, alkylthio, amino, alkylamino, or wherein R₂, R₃ and R₄ are a member of a ring system.

40 2. Compound according to claim 1, wherein R₂ is selected from the group consisting of hydrogen, mercapto and C₁-C₂₀-alkyl, preferably methyl, ethyl, propyl, isopropyl or butyl, halogen or amino.


45 3. Compound according to claim 1 or 2, wherein R₃ is hydrogen or is taken together with R₄ to form an, optionally substituted, alicyclic, aryl or heteroaryl ring system.

4. Compound according to claim 1, wherein R₁, R₂, R₃, R₄ and R₅ are independently selected from aryl, preferably phenyl and naphthyl, or heteroaryl, preferably furyl, pyrrolyl, thienyl, imidazolyl, thiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, benzothiazolyl, and oxadiazolyl.

50 5. Compound according to any of the preceding claims, wherein the compound is present in form of its addition salt, preferably hydrochloride, hydrobromide, phosphate, nitrate, acetate, malate, succinate, fumarate, tartrate, salicylate, sorbate, lactate, p-toluene, sulphate, or naphthalene-1,5-disulfonate salts.

55 6. Method for preparing a compound according to claim 1, comprising reacting a compound according to formula 2 with a compound according to formula 3 to prepare a compound according to formula 4, and reacting compound of formula 4 to result in compounds of formula 1, as given in scheme 1:

Scheme 1



with R_1 , R_2 , R_3 , R_4 and R_5 as defined in claim 1.

- 30
- 35
- 40
- 45
- 50
- 55
7. Method according to claim 6, wherein the reaction of compound 2 and compound 3 is in the presence of potassium carbonate in a solvent, preferably toluene, ethyl benzene, o-xylene- m-xylene, p-xylene, octane, nonane and/or isopropyl benzene.
 8. Method according to claim 6 or 7, wherein compound 4 is cyclized in the presence of at least one secondary amine, preferably diethyl amine, pyrrolidine, morpholine, piperidine and/or N-methylpiperazine, in a solvent, preferably toluene, ethylbenzene, o-xylene, m-xylene, p-xylene and/or isopropylbenzene.
 9. Pharmaceutical composition comprising at least one compound according to any of the claims 1-5 and a pharmaceutically acceptable carrier or excipient.
 10. Use of the compound according to any of the claims 1-5 or the pharmaceutical composition according to claim 9 as anticonvulsant agent for treatment of convulsion, especially for treatment of epilepsy.



EUROPEAN SEARCH REPORT

Application Number
EP 11 18 8961

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	H. I. EL-SUBBAGH ET AL.: "Synthesis and anticonvulsant activity of some new thiazolo[3,2-a][1,3]diazepine, benzo[d]thiazolo[5,2-a][12,6]diazepine and benzo[d]oxazolo[5,2-a][12,6]diazepine analogues", EUROPEAN JOURNAL OF MEDICINAL CHEMISTRY, vol. 46, no. 11, November 2011 (2011-11), pages 5567-5572, XP002670614, ISSN: 0223-5234, DOI: 10.1016/j.ejmech.2011.09.21 * page 5567, abstract; pages 5567, 5568, paragraph "Introduction"; page 5568, scheme 1, especially compounds 9 and 10; page 5569, table 1; page 5571, paragraphs 7.1.3. and 7.1.4. *	1-3,6-8, 10	INV. C07D513/04
X,D A	DE 103 20 732 A1 (FRIEDRICH SCHILLER UNI JENA BU [DE]) 9 December 2004 (2004-12-09) * claims 1,18; figure; example 2 *	6-8 1	
X,D A	EL-SUBBAGH ET AL: "New ultra-short acting hypnotic: Synthesis, biological evaluation, and metabolic profile of ethyl 8-oxo-5,6,7,8-tetrahydro-thiazolo[3,2-a][1,3]diazepin-3-carboxylate (HIE-124)", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, PERGAMON, ELSEVIER SCIENCE, GB, vol. 18, no. 1, 7 November 2007 (2007-11-07), pages 72-77, XP022410859, ISSN: 0960-894X, DOI: 10.1016/J.BMCL.2007.11.011 * page 73, scheme 1; page 74, figure 1 *	6-8 1	TECHNICAL FIELDS SEARCHED (IPC) C07D A61K A61P
----- -/--			
The present search report has been drawn up for all claims			
Place of search Berlin		Date of completion of the search 1 March 2012	Examiner Hass, Christian
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

1
EPO FORM 1503 03/02 (P04C01)



EUROPEAN SEARCH REPORT

Application Number
EP 11 18 8961

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	KADI ADNAN A ET AL: "Synthesis, ultra-short acting hypnotic activity, and metabolic profile of ethyl 8-oxo-5,6,7,8-tetrahydro-thiazolo[3,2-a][1,3]diazepin-3-carboxylate (HIE-124)", ARCHIV DER PHARMAZIE, WILEY - VCH VERLAG GMBH & CO. KGAA, DE, vol. 341, no. 2, 1 January 2008 (2008-01-01), pages 81-89, XP009148692, ISSN: 0365-6233, DOI: 10.1002/ARDP.200700132	6-8	
A	* page 82, scheme 1; page 83, figure 1 * -----	1	
A,D	QUARTARONE SILVANA ET AL: "Synthesis and anticonvulsant activity of N-3 substituted 2,3-benzodiazepines", FARMACO, SOCIETA CHIMICA ITALIANA, PAVIA, IT, vol. 59, 1 January 2004 (2004-01-01), pages 353-358, XP002489054, ISSN: 0014-827X, DOI: 10.1016/J.FARMAC.2004.01.005 * page 354, figure 1; page 357, scheme 1; page 357, table 1 * -----	1,10	
A,D	A. KADI ET AL.: "Liquid chromatographic high-throughput analysis of the new ultra-short acting hypnotic 'HIE-124' and its metabolite in mice serum using a monolithic silica column", ANALYST, vol. 136, no. 3, 7 February 2011 (2011-02-07), pages 591-597, XP002670615, * page 591, figure 1, compounds (A) and (B) * ----- -/--	1	TECHNICAL FIELDS SEARCHED (IPC)
The present search report has been drawn up for all claims			
Place of search Berlin		Date of completion of the search 1 March 2012	Examiner Hass, Christian
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

1
EPO FORM 1503 03/02 (P04C01)



EUROPEAN SEARCH REPORT

Application Number
EP 11 18 8961

DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim
A,D	<p>ROGAWSKI ET AL: "Therapeutic potential of excitatory amino acid antagonists: channel blockers and 2,3-benzodiazepines", TRENDS IN PHARMACOLOGICAL SCIENCES, ELSEVIER, HAYWARTH, GB, vol. 14, no. 9, 1 September 1993 (1993-09-01), pages 325-331, XP023843277, ISSN: 0165-6147, DOI: 10.1016/0165-6147(93)90005-5 [retrieved on 1993-09-01] * figure 2 *</p> <p style="text-align: center;">-----</p>	1
		CLASSIFICATION OF THE APPLICATION (IPC)
		TECHNICAL FIELDS SEARCHED (IPC)
The present search report has been drawn up for all claims		
1	Place of search Berlin	Date of completion of the search 1 March 2012
	Examiner Hass, Christian	
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>		

EPO FORM 1503 03.92 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 11 18 8961

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

01-03-2012

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 10320732	A1	09-12-2004	NONE

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- DE 10320732 [0005] [0020]

Non-patent literature cited in the description

- **BORDIE M.J.** *Lancet*, 1990, vol. 336, 350-354 [0002]
- **SHORAVON S.D.** *Lancet*, 1990, vol. 336, 93-96 [0002]
- In New Anticonvulsant Drugs. **MELDRUM B.S. et al.** Current Problems in Epilepsy. John Libby, 1986, vol. 4 [0002]
- **FLAHERTY P.T. et al.** *J. Med. Chem.*, 1996, vol. 39, 1509-1513 [0002]
- **DINGLELINE R. et al.** *Pharmacol. Rev.*, 1999, vol. 51, 7-61 [0004]
- **LEES G. J.** *Drugs*, 2000, vol. 59, 33-78 [0004]
- **ROGAWSKI M. A.** *Trends Pharmacol. Sci.*, 1993, vol. 14, 325-331 [0004]
- **QUARTARONE S. et al.** *IL Farmaco*, 2004, vol. 59, 353-358 [0004]
- **KADI et al.** *analyst*, 2011, vol. 136, 591-597 [0005]
- **MOLINA, P. et al.** *J. Org. Chem.*, 1993, vol. 58, 5264-5270 [0020]
- **IMMING, P. et al.** *Arch. Pharm.*, 1995, vol. 238, 207-215 [0020]
- **EI-SUBBAGH, H.I. et al.** *Bioorg. & Med. Chem. let.*, 2008, vol. 18, 72-77 [0020]
- **WHITE, H.S. et al.** *Antiepileptic Drugs*. Raven, 1995, 99-121 [0022] [0048] [0049] [0052]
- **DAUGE, V. et al.** *Neuropsychopharmacology*, 2001, vol. 25, 690-98 [0022] [0051]
- **GHOSH, M.** *Fundamental of Experimental Pharmacology*. Scientific Book Agency, 1984, 153-158187-189 [0023]
- **GHOSH; M.** *Fundamentals of Experimental Pharmacology*. Scientific Book Agency, 1984, 153-158187-189 [0053]
- **LÖSCHER, W. et al.** *Epilepsy Res.*, 1991, vol. 9, 1-10 [0054]